

Ab initio extracting of separate electronic contributions to transition metals' transport properties

Galtsov I.S.^{1,2,®}, Minakov D.V.¹ and Levashov P.R.¹

¹ Joint Institute for High Temperatures of the Russian Academy of Sciences, Izhorskaya 13 Bldg 2, Moscow, 125412, Russia

² Moscow Institute of Physics and Technology, Institutskiy Pereulok 9, Dolgoprudny, 141701, Russia

® galtsov.is@phystech.edu

The s - d scattering mechanism, first proposed by Mott [1], is the dominant source of electrical resistivity in transition metals, primarily driven by electron–phonon interactions. Accurate modeling of s - d electronic transitions is essential for understanding processes under ultrafast laser pulses in thin metallic films. Allen's approach to transport properties calculations [2] applies broadly across metals and successfully reproduces the unusual nonlinear temperature dependence of resistivity in several transition metals. It has been suggested [3] that the ρ_{ss} contribution—where both initial and final scattering states are s -like—in Pd can be approximated by the resistivity of Ag—the nearest non-transition-metal neighbor of Pd. In this work, we consider several criteria for distinguishing Bloch states of s - and d -type to more accurately select strongly hybridized states close to the chemical potential. We compute separate electronic contributions to total resistivity of Pd using Allen's method built upon *ab initio* calculations and Wannier interpolation as implemented in our SKiES code [2]. The results presented for both electrical resistivity and phonon linewidth in Pd.

The work was supported by the Russian Science Foundation (Grant No. 24-79-00136).

[1] Mott N F 1964 *Adv. Phys.* **13** 325–422

[2] Galtsov I S, Fokin V B, Minakov D V and Levashov P R 2025 *Comput. Phys. Commun.* **317** 109834

[3] Fradin F Y 1974 *Phys. Rev. Lett.* **33**(3) 158–161