

Study of photophysical characteristics of H3 centers in diamond

Pomazkin D.A.^{1,®}, Danilov P.A.¹, Vins V.G.¹,
Skorikov M.L.¹, Vyatkin S.V.² and Kudryashov S.I.¹

¹ Lebedev Physical Institute of the Russian Academy of Sciences, Leninsky Avenue 53, Moscow, 119991, Russia

² Lomonosov Moscow State University, Leninskiye Gory 1, Moscow, 119991, Russia

® dpomazkin@lebedev.ru

Diamond color centers are in high demand in emerging photonic technologies such as quantum computing, single-photon sources, and magnetic field sensors. H3 centers are of great interest as an active medium for laser generation in the 500-600 nm range [1] [2]. Implementing such technologies requires precise measurements of the photophysical characteristics of the centers, which are currently known from outdated measurements from the 1980s [3].

In this study, the emission cross section was measured using two methods. The first method is based on studying the photoluminescence yield under pumping with ultrashort 150-fs laser pulses of variable intensity. The second method determines the emission cross section based on the luminescence decay kinetics. The values obtained using these methods, $2.7 \times 10^{15} \text{ cm}^2$ and $2.0 \times 10^{-15} \text{ cm}^2$, agree well with each other and exceed the published value of $1.6 \times 10^{-16} \text{ cm}^2$ by an order of magnitude.

The study was supported by the Russian Science Foundation (Grant No. 25-42-02002).

- [1] Aharonovich I and Neu E 2014 *Advanced Optical Materials* **2** 911–928 ISSN 2195-1071 URL <http://dx.doi.org/10.1002/adom.201400189>
- [2] Mironov V P, Protasova E A, Lipatov E I and Martynovich E F 2021 *AIP Conference Proceedings* **2392** 030001 ISSN 0094-243X URL <https://doi.org/10.1063/5.0062002>
- [3] Rand S 1986 Synthetic diamond for color center lasers *Advanced Solid State Lasers* (Optica Publishing Group) p FA9 URL <https://opg.optica.org/abstract.cfm?URI=ASSL-1986-FA9>