

Exotics mixed phase appearance in calculation of spatial charge profiles of plasma

Chigvintsev A.Yu.^{1,®}, Iosilevskiy I.L.^{1,2}, Noginova I.Yu.³
and Zorina I.G.⁴

¹ Moscow Institute of Physics and Technology, Institutskiy Pereulok 9,
Dolgoprudny, 141701, Russia

² Joint Institute for High Temperatures of the Russian Academy of Sciences,
Izhorskaya 13 Bldg 2, Moscow, 125412, Russia

³ National University of Science and Technology MISIS, Leninskiy Avenue 4,
Moscow, 119049, None

⁴ Bauman Moscow State Technical University, 2nd Baumanskaya Street 5,
Moscow, 105005, Russia

® alex012008@gmail.com

The paper discusses the possibility of the appearance of discontinuities in the results of calculations of equilibrium space charge profiles in the vicinity of the source of inhomogeneity [1]. These discontinuities are considered as a kind of micro-level manifestation of phase transitions in the local equation of state (EOS), which is used to describe the non-ideal electronic and/or ionic subsystem within the framework of the quasi-homogeneity approximation (“local density”) [2]. Particular attention in this work is paid to the possibility of a specific manifestation of the above-mentioned nonideality effects in the studied equilibrium charge profiles in the form of an ultradisperse two-phase mixture (“mixed phase”). The proposed general conclusion is the statement that the concept of mixed phase is not an attribute of exclusively astrophysical applications, but is a fairly general property of computational schemes used to describe equilibrium inhomogeneous Coulomb systems [3].

[1] Iosilevski I, Chigvintsev A, Noginova L and Zorina I 2022 *High Temperature* **60** 325

[2] Iosilevski I 1985 *High Temperature* **23** 807 URL [arXiv:0901.3535](https://arxiv.org/abs/0901.3535).

[3] Iosilevski I, Chigvintsev A, Noginova L and Zorina I 2018 *J. of Phys. Conf. Ser.* **946** 012092