

Reconstruction of Z-pinch emission spectra in the wavelength range of less than 10 Å using a crystal X-ray spectrograph

Skobliakov A.V.^{1,®}, Kolesnikov D.S.¹, Kantsyrev A.V.¹
and Golubev A.A.¹

¹ National Research Center “Kurchatov Institute, Kurchatov Square 1,
Moscow, 123182, Russia

® dinAlt220@yandex.ru

This work is devoted to the reconstruction of Z-pinch plasma emission spectra in the wavelength range of less than 10 Å recorded by using a crystal x-ray spectrograph at the Angara 5-1 mega-ampere facility. The spectrograph JA-1 used in experiments has a cylindrical mica crystal with dimensions of $50 \times 40 \text{ mm}^2$ and radius of curvature of 100 mm. Registration of spectra is performed on the photographic film UF-4 with dimensions of $30 \times 10 \text{ mm}^2$. To reconstruct the spectra, the previously developed method based on iterative approximation of a true spectrum shape while minimizing a residual between experimental and calculated spectrograms is used [1,2]. The calculated spectrogram was obtained taking into account the instrumental function of the spectrograph. To define the instrumental function a virtual Monte-Carlo model in the Geant4 toolkit has been developed [3]. This model takes into account the interaction of radiation with the mica crystal using dynamical theory of diffraction. A true spectrum of Z-pinch plasma radiation is reconstructed for a load made of Al wire liners.

- [1] Skobliakov A V *et al.* 2023 *Plasma Physics Reports* **49** ISSN 1562-6938
- [2] Skobliakov A V *et al.* 2023 *Review of Scientific Instruments* **94** ISSN 0034-6748
- [3] Skobliakov A V *et al.* 2024 *Review of Scientific Instruments* **95** ISSN 0034-6748