

Electrical resistivity of liquid expanded metals by Ziman approach

Migdal K P^{1,2,®}, Petrov Yu V^{2,3}, Zhakhovsky V V^{1,2} and Inogamov N A^{1,2}

¹ Dukhov Research Institute of Automatics (VNIIA), Sushchevskaya 22, Moscow 127055, Russia

² Landau Institute for Theoretical Physics of the Russian Academy of Sciences, Akademika Semenova 1a, Chernogolovka, Moscow Region 142432, Russia

³ Moscow Institute of Physics and Technology, Institutskiy Pereulok 9, Dolgoprudny, Moscow Region 141700, Russia

® kir-migdal@yandex.ru

Transport properties of metals near the critical points were intensively studied [1, 2]. Using quantum dynamics simulations, we consider the structures of liquid aluminum and copper at high equilibrium temperatures and strong compression. On the other hand, we carry out classical molecular dynamics (MD) simulations using the embedded-atom method (EAM) interatomic potentials. Both approaches provide data for resistivity calculations based on the well-known Ziman–Evans (ZE) formula. The results of these calculations show a weak dependence on the used methods of ion structure computations. Also, we demonstrate that an effect of electron temperature, which is implemented in the effective electron–ion potential used in the ZE formula, is negligible at electron temperatures less than 30 000 K. This work has been supported by Russian Science Foundation (grant No. 14-19-01599).

[1] Clérrouin J, Noiret P, Korobenko V N and Rakhel A D 2008 *Phys. Rev. B* **78** 224203

[2] Khomkin A L and Shumikhin A S 2016 *JETP* **123** 891–902